

SALMON AND STEELHEAD HABITAT LIMITING FACTORS

**WATER RESOURCE INVENTORY AREA 17
QUILCENE-SNOW BASIN**



Photo provided by Hilton Turnbull

**WASHINGTON STATE
CONSERVATION COMMISSION**

FINAL REPORT

**Ginna Correa
November 2002**

ACKNOWLEDGEMENTS

The WRIA 17 salmon habitat limiting factors report could not have been completed without considerable contributions of time and effort from the following people who participated in various capacities on the technical advisory group (TAG):

Al Latham, Jefferson County Conservation District
Barbara Nightingale, MMA, Marine Resource Scientist, Biomes
Byron Rot, Jamestown S’Klallam Tribe
Chris May, PhD, Watershed Ecology LLC
Ed Manary, Washington Conservation Commission
Glenn Gately, Jefferson County Conservation District
Hilton Turnbull, Jamestown S’Klallam Tribe
John Cambalik, Puget Sound Action Team
Mark McHenry, US Forest Service
Randy Johnson, Washington Department of Fish and Wildlife
Richard Brocksmith, Hood Canal Coordinating Council
Steve Todd, Point No Point Treaty Council
Ted Labbe, Port Gamble S’Klallam Tribe
Thom Johnson, Washington Department of Fish and Wildlife

In addition, the author also wishes to thank the following for extensive information regarding fish populations and habitat conditions and substantial editorial comments during development of the report: Andy Perleberg, JSKT, for contribution to fish distribution; Dr. Carol Smith, WCC, for initial guidance on process and contribution to the Introduction chapter of this report; Dr. Charles Simenstad, UW for initial guidance on nearshore processes; Chris Byrnes, WDFW, for contribution to fish distribution; Hugh Shipman, Ecology, for initial guidance on nearshore processes; Ian Jablonski, City of Port Townsend, for water quantity information in the Big and Little Quilcene watersheds; Jeff Davis, WDFW, for participation in watershed review; Jennifer Cutler, NWIFC, for fish distribution mapping/table development and watershed/subbasin mapping; Larry Telles, USFWS, for participation in Big Quilcene review; Mike Kuttel, Jr., WCC, for extensive editing of this report and valuable computer assistance; Paula Mackrow, North Olympic Salmon Coalition, for participation in watershed and nearshore review; Peter Bahls, NWI, for contribution to watershed review; Randy Cooper, WDFW, for stock status and distribution; Ron McFarland, NWIFC, for fish distribution and shoreline modification mapping; and Tim Rymer, WDFW, contribution to fish distribution.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	2
TABLE OF CONTENTS	3
LIST OF TABLES	7
LIST OF FIGURES	8
LIST OF MAPS	12
ABBREVIATIONS AND ACRONYMS	13
EXECUTIVE SUMMARY	14
Introduction	18
Salmonid Habitat Limiting Factors Background	18
The Relative Role of Habitat in Healthy Populations of Natural Spawning Salmon (Chapter Author — Carol Smith, PhD, edited by the report Author)	19
Watershed Characterization	24
Overview	24
Marple/Jackson Creek – WRIA 17.0001	26
Spencer Creek – WRIA 17.0004	27
Indian George Creek – WRIA 17.0011	27
Big Quilcene River – WRIA 17.0012	27
Little Quilcene River – WRIA 17.0076	28
Donovan Creek – WRIA 17.0115	29
Lindsay Creek – WRIA 17.0123	29
Tarboo Creek – WRIA 17.0129	29
Camp Discovery Creek – WRIA 17.0141	30
Fisherman Harbor Creek – WRIA 17.0153	30
Thorndyke Creek – WRIA 17.0170	30
Nordstrom Creek – WRIA 17.0180	31
Shine Creek – WRIA 17.0181	31
Bones – WRIA 17.181A	31
Ludlow Creek – WRIA 17.0192	31
Pidling Creek – WRIA 17.0200	32
Little Goose Creek – WRIA 17.0200A	32
Chimacum Creek – WRIA 17.0203	32
Snow Creek – WRIA 17.0219	33
Salmon Creek Watershed – WRIA 17.0245	34
Contractors Creek – WRIA 17.0270	36
Eagle Creek – WRIA 17.0272	36
Chicken Coop Creek – WRIA 17.0278	37
Jimmycomelately Creek- WRIA 17.0285	37
Dean Creek – WRIA 17.0293	38
Johnson Creek – WRIA 17.0301	38
Nearshore	38
Distribution and condition of stocks	40
Chinook (<i>Oncorhynchus tshawytscha</i>)	40
Chum (<i>Oncorhynchus keta</i>)	41

Coho (<i>Oncorhynchus kisutch</i>)	48
Pink Salmon (<i>Oncorhynchus gorbuscha</i>)	52
Winter Steelhead (<i>Oncorhynchus mykiss</i>).....	52
Rainbow Trout (<i>Oncorhynchus mykiss</i>).....	54
Status Summary.....	54
Fish Distribution.....	54
HABITAT LIMITING FACTORS BY SUB-BASIN	57
Big Quilcene Sub-Basin.....	58
Marple/Jackson Creek.....	58
Spencer Creek	61
Indian George Creek	65
Big Quilcene River, mouth to Rogers Street, river mile 1.0	69
Big Quilcene River, Rogers Street Bridge to Hiddendale, river mile 1.0-3.2.....	73
Big Quilcene River, river mile 3.2 to the falls at river mile 7.8.....	77
Big Quilcene River, above the falls at river mile 7.8, including Tunnel Creek and Townsend Creek.....	79
Penny Creek, tributary to the Big Quilcene River at river mile 2.8.....	82
Little Quilcene Sub-Basin	85
Little Quilcene River – mouth to river mile 2.7	85
Little Quilcene River – river mile 2.7 to 6.8	89
Little Quilcene River – above river mile 6.8.....	92
Leland Creek	94
Ripley Creek.....	97
Howe Creek.....	100
Donovan/Jakeway - WRIA 17.0115/17.0116	103
Tarboo/Thorndyke Sub-Basin	107
Lindsay Creek	107
Tarboo Creek, mouth to river mile 0.9.....	110
Tarboo Creek, river mile 0.9 to 4.0	113
Tarboo Creek, above river mile 4.0.....	116
East Fork Tarboo Creek	118
Camp Discovery Creek	121
Fisherman Harbor Creek	123
Thorndyke Creek.....	125
Ludlow Sub-Basin.....	129
Nordstrom Creek	129
Shine Creek, mouth to SR 104.....	132
Shine Creek – Above SR 104.....	135
Bones Creek	138
Ludlow Creek – Mouth to River Mile 0.5 (Cascade).....	139
Ludlow Creek – above river mile 0.5.....	141
Piddling Creek.....	143
Little Goose Creek	145
Chimacum Subbasin.....	147
Chimacum Creek, mouth to river mile 3.0.....	148
Chimacum Creek, river mile 3.0-9.4.....	152

Chimacum Creek, above river mile 9.4.....	156
Chimacum Creek, East Fork	159
Putansuu Creek.....	163
Naylor Creek	166
South Fork Chimacum Creek (Barnhouse Creek).....	168
Discovery Bay Subbasin	173
Snow Creek, mouth to river mile 3.5	173
Snow Creek, above river mile 3.5	177
Andrews Creek, mouth to Bolton Road	179
Andrews Creek, above Bolton Road	182
Salmon Creek, mouth to Uncas Road	184
Salmon Creek, above Uncas Road	187
Contractor's Creek	190
Eagle Creek	193
Sequim Bay Subbasin.....	197
Chicken Coop Creek	197
Jimmycomelately Creek, mouth to river mile 1.0	200
Jimmycomelately Creek, river mile 1.0-3.5	203
Jimmycomelately Creek, above river mile 3.5.....	205
East Fork Jimmycomelately Creek	207
Dean Creek.....	210
Johnson Creek	212
Habitat Condition Rating Standards and WRIA 17 Habitat Ratings	216
Nearshore Introduction.....	226
Jackson Cove through Quilcene Bay	227
Right Smart Cove	227
Wawa Point	228
Jackson Cove/Pulali Point.....	229
Dabob Bay/Point Whitney/Dabob Bay	230
Quilcene Bay	232
Dabob Bay through Thorndyke Bay	237
Dabob Bay.....	237
Southern and Eastern Toandos Peninsula	241
Thorndyke Bay.....	243
Nordstrom Creek to Portage Bay	245
Nordstrom Creek to Bridgehaven	245
Squamish Harbor.....	247
Termination Point to Point Hannon	249
Point Hannon to Tala Point	252
Ludlow Bay to Basalt Point	253
Mats Mats to Olele Point.....	255
Olele Point to Portage Canal	256
Portage Canal to McCurdy Point	261
Portage Canal to Kinney Point.....	261
East and North Marrowstone Island.....	262
Kilisut Harbor.....	264

East Port Townsend Bay/North and West Indian Island.....	266
Portage Canal to Kala Point	267
Kala Point to Point Hudson.....	269
Point Hudson to Point Wilson.....	271
Point Wilson to McCurdy Point.....	273
Discovery Bay	276
Cape George.....	276
Becket Point	278
Adelma Beach to Salmon/Snow Creeks.....	279
Snow/Salmon Creek Estuaries	280
Mill Point Spit to Diamond Point.....	282
Diamond Point to Rocky Point.....	284
Protection Island.....	285
Sequim Bay	288
Rocky Point to Hardwick Point.....	288
Hardwick Point to Chicken Coop Creek	289
Chicken Coop, Jimmycomelately and Dean Creek Estuaries	290
Dean Creek to Pitship Point	291
Prioritized Nearshore Action Recommendations.....	294
Reference List	311
Appendix A: Maps	316

LIST OF TABLES

Table 1. Quilcene National Fish Hatchery Summer Chum Releases. Data provided by Larry Telles, USFWS.....	42
Table 2. WRIA 17 Fish Distribution.....	55
Table 3. Spencer Creek Water Temperatures. Data provided by Ted Labbe, PGST.....	64
Table 4. Lower Big Quilcene Water Temperatures. Data provided by Ted Labbe, PGST.....	76
Table 5. Lower Little Quilcene River Water Temperatures. Data provided by Port Gamble S'Klallam Tribe.....	88
Table 6. Upper Little Quilcene River Water Temperature. Data provided by Port Gamble S'Klallam Tribe.....	93
Table 7. Leland Creek Water Temperatures. Data provided by Port Gamble S'Klallam Tribe.....	96
Table 8. Leland Creek Dissolved Oxygen Concentrations. Data provided by Glenn Gately, JCCD.....	96
Table 9. Ripley Creek Water Temperature. Data collected by Port Gamble S'Klallam Tribe.....	99
Table 10. Howe Creek Water Temperature. Data provided by Port Gamble S'Klallam Tribe.....	101
Table 11. Donovan Creek Watershed Water Temperatures. Data provided by Port Gamble S'Klallam Tribe.....	105
Table 12. East Tarboo Creek Stream Temperatures. Data provided by Ted Labbe, PGST.....	120
Table 13. Thorndyke Water Temperatures. Data provided by Ted Labbe, PGST.....	127
Table 14. Nordstrom Creek Water Temperatures. Data provided by Ted Labbe, PGST.....	131
Table 15. Shine Creek Water Temperature. Data provided by Ted Labbe, PGST.....	134
Table 16. Lower Chimacum Creek Annual Instantaneous Maximum Temperatures, °C.....	150
Table 17. Middle Chimacum Creek Water Temperatures. Data provided by Glenn Gately, JCCD.....	155
Table 18. East Fork Chimacum Creek Water Temperatures. Data provided by Ted Labbe, PGST and Glenn Gately, JCCD.....	162
Table 19. Naylor Creek Water Temperatures. Data provided by Glenn Gately, JCCD.....	167
Table 20. Johnson Creek Water Temperatures Recorded Annual High. Data provided by Ed Chadd, Streamkeepers of Clallam County.....	214
Table 21. Salmonid Habitat Condition Rating Standards.....	217
Table 22. WRIA 17 Habitat Ratings.....	223
Table 23. Nearshore Stressors - Effects Table.....	296
Table 24. Nearshore Action Recommendations (Prioritized).....	297

LIST OF FIGURES

Figure 1. Summer Chum Spawning in Chimacum Creek. Photo provided by Al Latham, JCCD	41
Figure 2. Big Quilcene Summer Chum Escapement, 1968 to 2001. Data provided by Thom Johnson, WDFW.....	42
Figure 3. Little Quilcene Summer Chum Escapement, 1968 to 2001. Data provided by Thom Johnson, WDFW.....	43
Figure 4. Chimacum Creek Summer Chum Escapement, 1999 to 2001. Data provided by Randy Cooper, WDFW	43
Figure 5. Snow Creek Summer Chum Escapement, 1974 to 2001. Data provided by Thom Johnson, WDFW.....	44
Figure 6. Salmon Creek Summer Chum Escapement, 1974 to 2001. Data provided by Thom Johnson, WDFW.....	45
Figure 7. Jimmycomelately Summer Chum Escapement, 1982 to 2001. Data provided by Thom Johnson, WDFW.....	46
Figure 8. Big Quilcene Fall Chum Escapement, 1968 to 2001. Data provided by Thom Johnson, WDFW.....	46
Figure 9. Little Quilcene Fall Chum Escapement 1968 to 2001. Data provided by Thom Johnson, WDFW.....	47
Figure 10. Big Quilcene Fall Chum Hatchery Releases, 1992 to 2000. Data provided by Larry Telles, USFWS.....	47
Figure 11. Dabob Bay Coho Escapement 1968 to 2001. Data provided by Thom Johnson, WDFW.....	48
Figure 13. Chimacum Coho Cumulative Redd Counts, 1984 to 2001. Data provided by Randy Cooper, WDFW	49
Figure 14. Snow Creek Coho Escapement. 1968 to 2001. Data provided by Thom Johnson, WDFW	50
Figure 15. Snow Creek Coho Smolt Migration, 1986 to 2001. Data provided by Thom Johnson, WDFW	50
Figure 16. Jimmycomelately Coho Redd Counts, 1984 to 2001. Data provided by Randy Cooper, WDFW	51
Figure 17. Little Quilcene Winter Steelhead. Data provided by Thom Johnson	52
Figure 18. Discovery Bay Winter Steelhead Escapement, 1986 to 2001. Data provided by Thom Johnson.....	53
Figure 19. Marple/Jackson Watershed. Map provided by Jennifer Cutler, NWIFC.....	58
Figure 20. Spencer Creek Watershed. Map provided by Jennifer Cutler, NWIFC.....	62
Figure 21. Indian George Creek Watershed. Map provided by Jennifer Cutler, NWIFC	65
Figure 22. Indian George Creek Estuary, Pre-restoration. Graphic provided by Randy Johnson, WDFW	68
Figure 23. Indian George Creek Estuary, Post-restoration. Photo provided by Randy Johnson, WDFW	68
Figure 24. Big Quilcene Watershed, river mile 0.0 to 1.0. Map provided by Jennifer Cutler, NWIFC.....	69

Figure 25. Big Quilcene River Estuary. Graphics provided by Randy Johnson, WDFW	72
Figure 26. Big Quilcene Watershed, river mile 1.0 to 3.2. Map provided by Jennifer Cutler, NWIFC.....	73
Figure 27. Big Quilcene, Hiddendale to Falls. Map provided by Jennifer Cutler, NWIFC	77
Figure 28. Upper Big Quilcene Watershed. Map provided by Jennifer Cutler, NWIFC	79
Figure 29. Penny Creek Watershed. Map provided by Jennifer Cutler, NWIFC	82
Figure 30. Little Quilcene Watershed, river mile 0.0 to 2.7. Map provided by Jennifer Cutler, NWIFC.....	85
Figure 31. Little Quilcene River Estuary. Graphics provided by Randy Johnson, WDFW	89
Figure 32. Little Quilcene Watershed, river mile 2.7 to 6.8. Map provided by Jennifer Cutler, NWIFC.....	90
Figure 33. Upper Little Quilcene Watershed. Map provided by Jennifer Cutler, NWIFC	92
Figure 34. Leland Creek Watershed. Map provided by Jennifer Cutler, NWIFC	94
Figure 35. Ripley Creek Watershed. Map provided by Jennifer Cutler, NWIFC.....	97
Figure 36. Howe Creek Watershed. Map provided by Jennifer Cutler, NWIFC.....	100
Figure 37. Donovan/Jakeway Watershed. Map provided by Jennifer Cutler, NWIFC	103
Figure 38. Lindsay Creek Watershed. Map provided by Jennifer Cutler, NWIFC.....	107
Figure 39. Tarboo Creek, river mile 0.0 to 0.9. Map provided by Jennifer Cutler, NWIFC	110
Figure 40. Tarboo Creek, river mile 0.9 to 4.0. Map provided by Jennifer Cutler, NWIFC.....	113
Figure 41. Tarboo Creek, above river mile 4.0. Map provided by Jennifer Cutler, NWIFC	116
Figure 42. East Fork Tarboo Creek. Map provided by Jennifer Cutler, NWIFC.....	118
Figure 43. Camp Discovery Watershed. Map provided by Jennifer Cutler, NWIFC ...	122
Figure 44. Thorndyke Watershed. Map provided by Jennifer Cutler, NWIFC	125
Figure 45. Nordstrom Creek Watershed. Map provided by Jennifer Cutler, NWIFC.	129
Figure 46. Lower Shine Creek Watershed. Map provided by Jennifer Cutler, NWIFC	132
Figure 47. Upper Shine Creek Watershed. Map provided by Jennifer Cutler, NWIFC	136
Figure 48. Bones Creek Watershed. Map provided by Jennifer Cutler, NWIFC	138
Figure 49. Lower Ludlow Watershed. Map provided by Jennifer Cutler, NWIFC	139
Figure 50. Upper Ludlow Creek Watershed. Map provided by Jennifer Cutler, NWIFC	141
Figure 51. Piddling Creek Watershed. Map provided by Jennifer Cutler, NWIFC.....	143
Figure 52. Little Goose Creek Watershed. Map provided by Jennifer Cutler, NWIFC	145
Figure 53. Lower Chimacum Watershed, river mile 0.0 to 3.0. Map provided by Jennifer Cutler, NWIFC.....	148
Figure 54. Middle Chimacum Creek Watershed. Map provided by Jennifer Cutler, NWIFC	152

Figure 55. Chimacum Creek Habitat Restoration. Photo provided by Al Latham, JCCD	154
Figure 56. Upper Chimacum Creek Watershed. Map provided by Jennifer Cutler, NWIFC	156
Figure 57. East Fork Chimacum Creek Watershed. Map provided by Jennifer Cutler, NWIFC	159
Figure 58. Putaansuu Watershed. Map provided by Jennifer Cutler, NWIFC.....	163
Figure 59. Naylor Creek Watershed. Map provided by Jennifer Cutler, NWIFC	166
Figure 60. South Fork Chimacum (aka Barnhouse) Watershed. Map provided by Jennifer Cutler, NWIFC	168
Figure 61. Lower Snow Creek Watershed, mouth to river mile 3.5. Map provided by Jennifer Cutler, NWIFC	174
Figure 62. Upper Snow Creek, above river mile 3.5. Map provided by Jennifer Cutler, NWIFC	177
Figure 63. Lower Andrews Creek Watershed. Map provided by Jennifer Cutler, NWIFC	179
Figure 64. Andrews Creek above Bolton Road. Map provided by Jennifer Cutler, NWIFC	182
Figure 65. Lower Salmon Creek Watershed. Map provided by Jennifer Cutler, NWIFC	184
Figure 66. Upper Salmon Creek Watershed. Map provided by Jennifer Cutler, NWIFC	187
Figure 67. Contractor's Creek Watershed. Map provided by Jennifer Cutler, NWIFC.....	190
Figure 68. Eagle Creek Watershed. Map provided by Jennifer Cutler, NWIFC	193
Figure 69. Eagle Creek Estuary. Photo provided by Hilton Turnbull, JSKT.....	195
Figure 70. Chicken Coop Creek Watershed. Map provided by Jennifer Cutler, NWIFC	197
Figure 71. Lower Jimmycomelately Watershed. Map provided by Jennifer Cutler, NWIFC	200
Figure 72. Jimmycomelately Watershed, river mile 1.0 to 3.5. Map provided by Jennifer Cutler, NWIFC	203
Figure 73. Upper Jimmycomelately Watershed. Map provided by Jennifer Cutler, NWIFC	205
Figure 74. East Fork Jimmycomelately Watershed. Map provided by Jennifer Cutler, NWIFC	207
Figure 75. Dean Creek Watershed. Map provided by Jennifer Cutler, NWIFC	210
Figure 76. Johnson Creek Watershed. Map provided by Jennifer Cutler, NWIFC	212
Figure 77. Big Quilcene and Little Quilcene Estuary Modification. Graphics provided by Randy Johnson, WDFW	234
Figure 78. Tarboo Bay. 2001 Ecology Nearshore Oblique Photo	237
Figure 79. Naturally Eroding Bluff. 2001 Ecology Nearshore Oblique Photo	238
Figure 80. Thorndyke Estuary. 2001 Ecology Nearshore Oblique Photos	243
Figure 81. South Point Ferry and Salt Marsh. 2001 Ecology Nearshore Oblique Photo	245
Figure 82. Bridgehaven. 2001 Ecology Nearshore Oblique Photo	246
Figure 83. Oak Bay. Graphics provided by Randy Johnson, WDFW	259

Figure 84. Chimacum Creek Nearshore, Existing. Photo provided by Randy Johnson, WDFW	268
Figure 85. Chimacum Creek Nearshore, After Proposed Restoration. Graphic provided by Randy Johnson, WDFW.....	268
Figure 86. Pier at Fort Worden. Ecology 2001 Oblique Photo	272
Figure 87. Cape George Marina. Ecology 2001 Oblique Photo	277
Figure 88. Beckett Point. 2001 Ecology Nearshore Oblique Photo.....	278
Figure 89. Salmon and Snow Estuaries with Historic Overlay. Graphic by Randy Johnson, WDFW	281
Figure 90. Eagle Creek Estuary. Photo provided by Hilton Turnbull.....	283
Figure 91. Protection Island Marina. 2001 Ecology Nearshore Oblique Photo.....	286
Figure 92. Jimmycomelately Estuary with Proposed Restoration Actions. Graphic by Randy Johnson, WDFW.....	291

LIST OF MAPS

- [Map 1.](#) WRIA/Subbasin Overview
- [Map 2.](#) Chinook Distribution
- [Map 3.](#) Chum Distribution
- [Map 4.](#) Coho Distribution
- [Map 5.](#) Pink Distribution
- [Map 6.](#) Steelhead/Rainbow Distribution
- [Map 7.](#) Cutthroat Distribution
- [Map 8.](#) Fish Passage Barriers
- [Map 9.](#) Subbasin Segments
- [Map 10.](#) Big and Little Quilcene Subbasin Hydromodifications
- [Map 11.](#) Tarboo/Thorndyke Subbasin Hydromodifications
- [Map 12.](#) Shine/Ludlow Subbasin Hydromodifications
- [Map 13.](#) Chimacum Subbasin Hydromodifications
- [Map 14.](#) Discovery Bay Subbasin Hydromodifications
- [Map 15.](#) Sequim Bay Subbasin Hydromodifications
- [Map 16.](#) Big and Little Quilcene Subbasins, Jackson Cove through Quilcene Bay Driftcells
- [Map 17.](#) Tarboo/Thorndyke Subbasin, Dabob Bay/Toandos through Thorndyke Bay Driftcells
- [Map 18.](#) Shine/Ludlow Subbasin, Thorndyke Bay to Portage Canal Driftcells
- [Map 19.](#) Chimacum Subbasin, Portage Canal to Point Wilson, including Marrowstone and Indian Islands Driftcells
- [Map 20.](#) Discovery Bay Subbasin, Point Wilson through Discovery Bay, including Protection Island Driftcells
- [Map 21.](#) Sequim Bay Subbasin, Rocky Point through Sequim Bay Driftcells
- [Map 22.](#) Big and Little Quilcene Subbasins, Jackson Cove through Quilcene Bay Shoreline Alterations
- [Map 23.](#) Tarboo/Thorndyke Subbasin, Dabob Bay/Toandos through Thorndyke Bay Shoreline Alterations
- [Map 24.](#) Shine/Ludlow Subbasin, Thorndyke Bay to Portage Canal Shoreline Alterations
- [Map 25.](#) Chimacum Subbasin, Portage Canal to Point Wilson, including Marrowstone and Indian Islands Shoreline Alterations
- [Map 26.](#) Discovery Bay Subbasin, Point Wilson through Discovery Bay, including Protection Island Shoreline Alterations
- [Map 27.](#) Sequim Bay Subbasin, Rocky Point through Sequim Bay Shoreline Alterations

ABBREVIATIONS AND ACRONYMS

AIMT	Annual Instantaneous Maximum Temperature
7-DADMT	7-day Average of Daily Maximum Temperature
21-DADT	21-day Average Daily Temperature
BIBI	Benthic Invertebrate Biotic Index
CCCD	Clallam County Conservation District
CREP	Conservation Reserve Enhancement Program
Ecology	Washington State Department of Ecology
HCSEG	Hood Canal Salmon Enhancement Group
JCCD	Jefferson County Conservation District
JSKT	Jamestown S’Klallam Tribe
LWD	Large Woody Debris
Mg/L	Milligrams per Liter
NOSC	North Olympic Salmon Coalition
NTU	Nephelometric Turbidity Unit
NWIFC	Northwest Indian Fisheries Commission
PGST	Port Gamble S’Klallam Tribe
PNPTC	Point No Point Treaty Council
RM	River Mile
SaSI	Salmon and Steelhead Inventory
SASSI	Salmon and Steelhead Stock Inventory
SSHEAR	Salmon Screening, Habitat Enhancement and Restoration
SSHIAP	Salmon and Steelhead Habitat Inventory Assessment Project
TAG	Technical Advisory Group
TFW	Timber, Fish and Wildlife
USFS	United States Forest Service
USFWS	United States Fish and Wildlife Service
WCC	Washington Conservation Commission
WDFW	Washington Department of Fish and Wildlife
WDNR	Washington Department of Natural Resources
WOS	Wild Olympic Salmon
WRIA	Water Resource Inventory Area

EXECUTIVE SUMMARY

Water Resource Inventory Area (WRIA) 17 is located along the northeast corner of the Olympic Peninsula in western Washington State. The WRIA extends from the Marple/Jackson watershed in southeast Jefferson County northward and westward to, and including, the Johnson Creek watershed along the west side of Sequim Bay. It is bordered to the north by the Strait of Juan de Fuca, to the east by Admiralty Inlet, northern Puget Sound and Hood Canal, and to the south and west by the Olympic Mountains and associated foothills and floodplains. The majority of the WRIA 17 watersheds are small lowland drainages with headwaters in the low foothills of the Olympic Mountains. The WRIA lies within the Olympic “rainshadow”, which indicates a moderate climate with rainfall limited to 15 to 20 inches in Port Townsend and increasing to 70 to 80 inches along the foothills of eastern Olympic mountain range. The WRIA area covers approximately 626 square miles (401,000 acres) and includes portions of Jefferson and Clallam counties. Over seventy percent of the WRIA is privately owned with a population exceeding 23,000 people (Parametrix, Inc. 2000).

The majority of WRIA 17 watersheds provide spawning and rearing habitats for four species of salmon: coho, chum, steelhead, and searun cutthroat. The estuarine and nearshore habitats provide a critical migration corridor for juvenile salmon of all species. Both Hood Canal summer chum and Puget Sound chinook are federally listed as threatened under the Endangered Species Act. Summer chum are documented in many WRIA 17 streams: Big Quilcene River, Little Quilcene River, Chimacum Creek, Snow Creek, Salmon Creek and Jimmycomelately Creek. It has not been determined if native chinook still exist in Hood Canal due to the interrelations with a mixed composition of hatchery introductions. Artificial supplementation programs have contributed to inconsistent adult chinook returns to the Big Quilcene River and more recently to Tarboo Creek. Fall chum returning to Quilcene Bay and Dabob Bay watersheds are considered healthy. Coho stocks range from critical in Discovery Bay to depressed in Quilcene, Dabob and Sequim Bay watersheds. Chimacum coho are healthy according to the proposed 2002 Salmon and Steelhead Inventory (SaSI) but that designation is subject to debate due to the index location in the best spawning habitat. Not much is known about winter steelhead populations with the exception of the Discovery Bay stocks, which have been designated depressed by proposed SaSI.

The species found in WRIA 17 utilize specialized habitats at different times and at different life stages. Different species stagger their upstream migration and each species has a unique rearing strategy. All species require adequate flow and water quality, ample spawning gravels, instream structure in the form of large woody debris and/or large boulders, pools, and a functional riparian zone while inhabiting the riverine system. While coho, chinook and steelhead remain in the freshwater for an extended period of time following fry emergence, pink and chum salmon tend to move directly out into the salt water. Estuarine, high salt marsh, eelgrass and shallow water nearshore habitats are critical to all species of juvenile salmonids as they enter the marine environment. Pink and chum salmon rely heavily on eelgrass beds for feeding and hiding and shallow water

for prey avoidance. Studies also show that high salt marsh and estuarine tidal channels are critical habitats for chinook and coho as well.

Human alterations to salmonid habitat can be expected to have different consequences for different fish species and their life stages. While natural environmental conditions, such as fire, floods and mass wasting events create a disturbance/rebuilding cycle that tends to nourish the aquatic environment, human alterations to the landscape can impact the environment beyond its natural ability to heal and sustain fish resources. Freshwater rearing is particularly critical for coho whose typical freshwater cycle extends through the summer months when many WRIA 17 streams are flow-limited. Freshwater rearing salmonids are therefore particularly vulnerable to habitat impacts such as elevated water temperatures and dewatering as a result of riparian removal and water extraction, and lack of instream structure such as pool-forming large woody debris. In the marine environment, shoreline alterations, such as bank armoring, overwater structures and intertidal fill, can interrupt important sediment input from eroding bluffs, alongshore sediment movement or drift, and continuous eelgrass beds that are critical to migrating juvenile salmon.

The Habitat Limiting Factors Analysis for WRIA 17 summarizes existing salmonid habitat data and represents the most current compilation and review of riverine and nearshore processes and human-induced impacts to salmon productivity. It does not cover salmonid productivity limited by hydroelectric dams, harvest or hatcheries. Data included or referenced in this report include watershed analysis, formal habitat inventories or studies specifically directed at evaluating fish habitat, salmon stock inventories and assessments, and other watershed data not specifically associated with fish habitat evaluation. Where data are lacking, the Technical Advisory Group (TAG) relied on its combined professional knowledge to assess the extent to which habitat conditions are affecting salmonid productivity. Where data and best professional knowledge are lacking, the habitat elements have been identified as data gaps and warrant additional specific watershed research or evaluation.

Land use activities associated with forest practices, agriculture, rural development and shoreline development have had negative impacts on salmon habitat in WRIA 17. Habitat conditions in the federally owned lands in the upper watershed, managed by the US Park Service (USPS) and the US Forest Service (USFS), are among the best in the WRIA. The USPS strives to maintain natural habitats through preservation, and even though very little land in WRIA 17 is managed by the USPS, their conservation measures protect downstream riverine function. The USFS has adopted a Riparian Reserve Program which provides for well functioning riparian habitat that ensures conifer canopy cover for temperature control, large woody debris recruitment, streambank stability to limit fine sediment input, and migratory corridors for numerous wildlife species. Their clearcuts rarely exceed 80 acres in size. This is in contrast to the large clearcuts and inadequate riparian zones on state-owned and private forest lands as evidenced in orthophoto analysis by TAG members. The riparian zone's ability to intercept fine sediments resulting from exposed soils diminishes as the riparian buffers decrease. In addition, mass wasting events and the subsequent above-normal delivery of sediments

into the Snow Creek, Nordstrom Creek and Chimacum Creek watersheds have been directly linked to improper forest road construction, maintenance and/or abandonment on state-managed and/or private forest lands. Elevated water temperatures, lack of large woody debris, and limited large woody debris recruitment due to limited riparian areas are typical of many of the watersheds in the WRIA.

Agriculture activities within the floodplains of many WRIA 17 watersheds have channelized mainstems and tributaries, drained beaver ponds for livestock grazing and peat mining and eliminated forested riparian zones. These activities have eliminated valuable juvenile overwintering and rearing habitat associated with beaver ponds, decreased broad channel meanders, eliminated floodplain connectivity to side channel habitats, reduced channel complexity and instream structure, minimized pool/riffle ratios, decreased streambed and streambank stability, and eliminated healthy riparian zones. Salmon Creek, Chimacum Creek, Upper Ludlow Creek, and Tarboo Creek watersheds have experienced degradation from agriculture activities. Further agricultural impacts to estuarine and salt marsh habitats can be observed in the Big Quilcene, Little Quilcene and Snow Creek watersheds, where extensive diking and filling into the nearshore environment have decreased estuarine function and eliminated salt marsh habitats.

Jefferson County has designated Port Townsend as its urban growth area as defined by the Washington State Growth Management Act. This has not entirely eliminated rural development throughout the county where the total population growth is among the highest in the state. Rural development tends to remove riparian vegetation for views, increase stormwater runoff with impervious surfaces and threaten water quality through the use of herbicides, pesticides and chemical fertilizers. Shoreline development contributes further to impacts to salmonid habitats. Removal of riparian vegetation for views weakens bank stability which could threaten home sites and which often results in bank protection. Shoreline armoring prevents the movement of sediments from naturally occurring backshore feeder bluffs to the nearshore. Intertidal fill eliminates shallow water habitat utilized by migrating juvenile salmon to escape prey, fragments eelgrass beds, interrupts alongshore sediment drift, and can decrease valuable salt marsh and lagoon habitats.

In order to ensure that salmonid habitats can produce sustainable and harvestable populations into the future, the Technical Advisory Group consistently placed preservation of properly functioning habitats, particularly estuaries, actively eroding feeder bluffs and riverine riparian corridors, as priority action recommendations. Preservation of critical habitats is a cost effective tool to ensure that properly functioning habitats will remain as such into perpetuity. Protection of critical habitats is the top priority for Salmon Creek and the Big Quilcene River, and is an important action recommendation for Snow Creek and Chimacum Creek. The Chumsortium, a partnership between Jefferson Land Trust, Wild Olympic Salmon, Jefferson County Conservation District, North Olympic Salmon Coalition, Hood Canal Coordinating Council, Trout Unlimited, Jefferson County and the Washington Department of Fish and Wildlife, has been actively and strategically pursuing funding for the acquisition of critical salmon habitats in east Jefferson County.

When a watershed has been severely impacted and cannot heal itself within a reasonable time frame, habitat restoration may be necessary. Once the source of the problem has been identified, rehabilitation activities can be directed to restore properly functioning condition. Such activities in the riverine environment might include removal of artificial barriers to fish passage, reestablishment of a healthy riparian zone, restoration of sinuosity and/or complexity, installation of cattle exclusion fences, abatement of mass wasting events, and/or removal of streambank armoring. Restoration activities in the nearshore might include removal of intertidal fill, restoration of lagoon and/or salt marsh connectivity, removal of shoreline armoring and/or removal of estuary constrictions that impede natural function. In some cases, property acquisition may be necessary prior to initiating restoration activities. The Technical Advisory Group identified restoration activities for the majority of the watersheds in WRIA 17. Jefferson County Conservation District, Wild Olympic Salmon, North Olympic Salmon Coalition, Hood Canal Salmon Enhancement Group, Port Gamble S'Klallam Tribe, Jefferson County and Washington Department of Fish and Wildlife have all pursued funding and participated in habitat restoration projects. Continued efforts will be necessary to increase habitat productivity to ensure sustainable salmonid populations.

Protection and restoration activities are only a part of the salmonid habitat equation. Land use regulations and their enforcement must be redirected to protect the valuable fish and wildlife resources that WRIA 17 has to offer. Preventing habitat degradation is a very cost effective tool to ensure sustainable populations of fish and wildlife into the future. The Technical Advisory Group identified assessments and studies needed to fill data gaps. In some cases, assessments might be necessary prior to beginning preservation or restoration activities.

The following report is a detailed assessment of habitat limiting factors in WRIA 17. Each watershed assessment is complete with a prioritized list of action recommendations for that watershed. The nearshore discussion is followed by a prioritized list of nearshore projects for the entire WRIA. This report provides information that can be used in the development of salmonid habitat protection and restoration strategies. It is a snapshot in time that can be supplemented with additional data from habitat assessments and habitat restoration successes as information becomes available.